

Anomalous Transport from Fluid/Gravity Correspondence

Michael Lublinsky

Ben-Gurion University of the Negev, Beer Sheva, Israel

Yanyan Bu, M.L., Amir Sharon, 1608.08595 (JHEP), 1609.09054

Yanyan Bu and M.L., 1406.7222 (PRD), 1409.3095 (JHEP),
1502.08044 (JHEP)

Yanyan Bu, M.L., Amir Sharon, 1504.01370 (JHEP), 1511.08789 (JHEP)

Warm up: Electric Current

$$\vec{J} = -\mathcal{D}^0 \vec{\nabla} \rho + \sigma_e^0 \vec{\mathcal{E}}; \quad \vec{J}(\omega, \vec{q}) = -\mathcal{D}^0 i\vec{q} \rho(\omega, \vec{q}) + \sigma_e^0 \vec{\mathcal{E}}(\omega, \vec{q})$$

\mathcal{D}^0 is a diffusion constant; σ_e^0 is a DC conductivity.

Continuity equation (charge conservation): $\vec{\nabla} \cdot \vec{J} + \dot{\rho} = 0$

Linear response theory:

$$\vec{J}(\omega, \vec{q}) = \sigma_e^{\text{AC}}(\omega) \vec{\mathcal{E}}(\omega, \vec{q}); \quad \sigma_e^{\text{AC}} = \frac{i\omega \sigma_e^0}{i\omega - \mathcal{D}^0 q^2}$$

AC conductivity. Valid in hydrodynamic regime of small ω, q .

Any generalisations to finite ω and q ?

The most general (linear) constitutive relation for e/m current including both longitudinal and transverse responses

$$\vec{\mathbf{J}}(\omega, \vec{\mathbf{q}}) = -\mathcal{D}(\omega, \mathbf{q}^2) \mathbf{i}\vec{\mathbf{q}} \rho(\omega, \vec{\mathbf{q}}) + \sigma_e(\omega, \mathbf{q}^2) \vec{\mathcal{E}}(\omega, \vec{\mathbf{q}}) + \sigma_m(\omega, \mathbf{q}^2) \mathbf{i}\vec{\mathbf{q}} \times \vec{\mathcal{B}}(\omega, \vec{\mathbf{q}}) .$$

$$\vec{\mathbf{J}} = -\mathcal{D}(\partial^t, \nabla^2) \vec{\nabla} \rho + \sigma_e(\partial^t, \nabla^2) \vec{\mathcal{E}} + \sigma_m(\partial^t, \nabla^2) \vec{\nabla} \times \vec{\mathcal{B}}.$$


This is an "off-shell" constitutive relation.

\mathcal{D} , σ_e , and σ_m are momenta-dependent transport coefficient functions (TCF).

can be uniquely determined in holographic models to be discussed next

AdS/CFT

AdS/QCD



*Changes in
length scale
mapped to
evolution in the
5th dimension **z***

Maxwell field in Schwarzschild- AdS_5 geometry (probe approximation)

$$S = - \int d^5x \sqrt{-g} \frac{1}{4} e^2 (F^V)_{MN} (F^V)^{MN} + S_{\text{c.t.}}$$

Maxwell equations

$$EQ^N := \nabla_M F^{MN} = 0$$

Schwarzschild- AdS_5 geometry (ingoing Eddington-Finkelstein coordinates)

$$ds^2 = g_{MN} dx^M dx^N = 2dt dr - r^2 f(r) dt^2 + r^2 \delta_{ij} dx^i dx^j,$$

$f(r) = 1 - 1/r^4$. The horizon is at $r = 1$, the Hawking temperature is $\pi T = 1$.

Near the conformal boundary $r = \infty$ the solution is expandable in a series ($A_r = 0$)

$$A_\mu(r, x_\alpha) = A_\mu^{(0)}(x_\alpha) + \frac{A_\mu^{(1)}(x_\alpha)}{r} + \frac{A_\mu^{(2)}(x_\alpha)}{r^2} + \frac{B_\mu^{(2)}(x_\alpha)}{r^2} \log r^{-2} + \mathcal{O}\left(\frac{\log r^{-2}}{r^3}\right),$$

The boundary current (using the holographic dictionary)

$$J^\mu = -\eta^{\mu\nu} \left(2A_\nu^{(2)} + 2B_\nu^{(2)} + \eta^{\sigma t} \partial_\sigma F_{t\nu}^{(0)} \right).$$

4 dynamical eqns $\mathbf{EQ}^\mu = 0 \rightarrow$ transport, $\mathbf{EQ}^r = 0 \rightarrow$ current conservations.

$\mathbf{EQ}^\mu = 0$ admit the most general static homogeneous solutions

$$\mathbf{A}_\mu = \mathbf{A}_\mu^{(0)} + \frac{\rho}{2r^2} \delta_{\mu t}, \quad \mathbf{A}_\mu^{(0)} = \text{const}, \quad \rho = \text{const}$$

The boundary theory is a static uniformly charged plasma with no external fields

$$\mathbf{J}^t = \rho, \quad \mathbf{J}^i = 0$$

Next, following the spirit of

S. Bhattacharyya, V. E. Hubeny, S. Minwalla, and M. Rangamani, (2008)

$$\mathbf{A}_\mu^{(0)} \rightarrow \mathbf{A}_\mu^{(0)}(\mathbf{x}_\alpha), \quad \rho \rightarrow \rho(\mathbf{x}_\alpha).$$

The solution has to be amended:

$$\mathbf{A}_\mu(\mathbf{r}, \mathbf{x}_\alpha) = \mathbf{A}_\mu^{(0)}(\mathbf{x}_\alpha) + \frac{\rho(\mathbf{x}_\alpha)}{2r^2} \delta_{\mu t} + \mathbf{a}_\mu(\mathbf{r}, \mathbf{x}_\alpha)$$

Solve for a (bulk-to-boundary propagator)

Different from approaches based on two-point correlators, which assume on-shellness

$U(1)$ vector current: Diffusion and Conductivity

$$\vec{\mathbf{J}}(\omega, \vec{\mathbf{q}}) = -\mathcal{D}(\omega, \mathbf{q}^2) \mathbf{i}\vec{\mathbf{q}} \rho(\omega, \vec{\mathbf{q}}) + \sigma_e(\omega, \mathbf{q}^2) \vec{\mathbf{E}}(\omega, \vec{\mathbf{q}}) + \sigma_m(\omega, \mathbf{q}^2) \mathbf{i}\vec{\mathbf{q}} \times \vec{\mathbf{B}}(\omega, \vec{\mathbf{q}}).$$

$$\mathcal{D} = \frac{1}{2} + \frac{1}{8}\pi\mathbf{i}\omega + \frac{1}{48} \left[-\pi^2\omega^2 + \mathbf{q}^2 (6 \log 2 - 3\pi) \right] + \dots,$$

$$\sigma_e = 1 + \frac{\log 2}{2}\mathbf{i}\omega + \frac{1}{24} \left[\pi^2\omega^2 - \mathbf{q}^2 (3\pi + 6 \log 2) \right] + \dots,$$





$$\sigma_m = 0 + \frac{1}{16}\mathbf{i}\omega \left(2\pi - \pi^2 + 4 \log 2 \right) + \dots.$$

$\sigma_m^0 > 0$ in a pure QED plasma with one Dirac fermion at one loop level

B. B. Brandt, A. Francis, and H. B. Meyer, (2014)



$\sigma_m^0 = 0$ based on Boltzmann equations J. Hong and D. Teaney, (2010)





Maxwell is linear (exact), no Lorentz force



Memory Function / Causality

$$\tilde{\mathcal{D}}(t, q^2) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \mathcal{D}(\omega, q^2) e^{-i\omega t} d\omega$$





$U_V(1) \times U_A(1)$: Anomaly-induced transport

5d Lagrangian:

$$\mathcal{L} = -\frac{1}{4}e^2(\mathbf{F}^V)_{MN}(\mathbf{F}^V)^{MN} - \frac{1}{4}e'^2(\mathbf{F}^a)_{MN}(\mathbf{F}^a)^{MN} \\ + \frac{\kappa \epsilon^{MNPQR}}{2\sqrt{-g}} \left[3e^2 e' \mathbf{A}_M (\mathbf{F}^V)_{NP} (\mathbf{F}^V)_{QR} + e'^3 \mathbf{A}_M (\mathbf{F}^a)_{NP} (\mathbf{F}^a)_{QR} \right].$$

Boundary currents:

$$\partial_\mu \mathbf{J}^\mu = 0, \quad \partial_\mu \mathbf{J}_5^\mu = 4\kappa \left(3\vec{\mathcal{E}} \cdot \vec{\mathcal{B}} + \vec{\mathcal{E}}^a \cdot \vec{\mathcal{B}}^a \right)$$

external e/m $(\vec{\mathcal{E}}, \vec{\mathcal{B}})$ and axial $(\vec{\mathcal{E}}^a, \vec{\mathcal{B}}^a)$ fields.

CS introduces non-linearity in EQ

I) Linear transport

$$\rho(\mathbf{x}_\alpha) = \bar{\rho} + \epsilon \delta \rho(\mathbf{x}_\alpha), \quad \rho_5(\mathbf{x}_\alpha) = \bar{\rho}_5 + \epsilon \delta \rho_5(\mathbf{x}_\alpha),$$

$$\mu(\mathbf{x}_\alpha) = \bar{\mu} + \epsilon \delta \mu(\mathbf{x}_\alpha), \quad \mu_5(\mathbf{x}_\alpha) = \bar{\mu}_5 + \epsilon \delta \mu_5(\mathbf{x}_\alpha), \quad \bar{\mu} = \bar{\rho}/2, \quad \bar{\mu}_5 = \bar{\rho}_5/2$$

$$\mathcal{E}_i(\mathbf{x}_\alpha) \rightarrow \epsilon \mathcal{E}_i(\mathbf{x}_\alpha), \quad \mathcal{B}_i(\mathbf{x}_\alpha) \rightarrow \epsilon \mathcal{B}_i(\mathbf{x}_\alpha), \quad \mathcal{E}_i^a(\mathbf{x}_\alpha) \rightarrow \epsilon \mathcal{E}_i^a(\mathbf{x}_\alpha), \quad \mathcal{B}_i(\mathbf{x}_\alpha) \rightarrow \epsilon \mathcal{B}_i^a(\mathbf{x}_\alpha).$$

$$\mathbf{J}^t = \rho, \quad \vec{\mathbf{J}} = -\mathcal{D} \vec{\nabla} \rho + \sigma_e \vec{\mathcal{E}} + \sigma_m \vec{\nabla} \times \vec{\mathcal{B}} + \sigma_\chi \vec{\mathcal{B}} + \sigma_a \vec{\nabla} \times \vec{\mathbf{B}}^a + \sigma_\kappa \vec{\mathcal{B}}^a$$

$$\mathbf{J}_5^t = \rho_5, \quad \mathbf{J}_5^i = -\mathcal{D} \vec{\nabla} \rho_5 + \sigma_e \vec{\mathcal{E}}^a + \sigma_m \vec{\nabla} \times \vec{\mathcal{B}}^a + \sigma_\chi \vec{\mathcal{B}}^a + \sigma_a \vec{\nabla} \times \vec{\mathcal{B}} + \sigma_\kappa \vec{\mathcal{B}}. \quad \partial_\mu \mathbf{J}_5^\mu = 0$$

σ_χ – **CME**; **D. E. Kharzeev and H. J. Warringa, (2009)**

σ_κ – **CSE**; **D. T. Son and A. R. Zhitnitsky, (2004) ; M. A. Metlitski and A. R. Zhitnitsky, (2005)**

$$\sigma_{\text{m}} = 72\kappa^2 \left(\bar{\mu}^2 + \bar{\mu}_5^2 \right) (2 \log 2 - 1) + \text{i}\omega \left[\frac{1}{16}(2\pi - \pi^2 + 4 \log 2) + \mathcal{O} \left(\bar{\mu}^2 + \bar{\mu}_5^2 \right) \right] + \cdots,$$

$$\sigma_{\text{m}}[\mathbf{q} = \mathbf{0}] - \sigma_{\text{m}}[\mathbf{q} = \mathbf{0}, \bar{\mu} = \bar{\mu}_5 = 0] \text{ is linear in } \kappa^2 (\bar{\mu}^2 + \bar{\mu}_5^2)$$

$$\sigma_{\chi} = 12\kappa\bar{\mu}_5 \left\{ 1 + \text{i}\omega \log 2 - \frac{1}{4}\omega^2 \log^2 2 - \frac{q^2}{24} \left[\pi^2 - 1728\kappa^2 \left(\bar{\mu}_5^2 + 3\bar{\mu}^2 \right) (\log 2 - 1)^2 \right] \right\} + \cdots,$$


$$\sigma_{\chi}^0 \quad \text{A. Gynther, K. Landsteiner, F. Pena-Benitez, and A. Rebhan, (2011)}$$

$$\sigma_{\chi}[\mathbf{q} = \mathbf{0}] \text{ is linear in } \kappa \mu_5 \text{ and independent of } \mu$$


$$\sigma_{\kappa}[\mu, \mu_5] = \sigma_{\chi}[\mu_5, \mu] \qquad \sigma_a = 144\kappa^2 \bar{\mu} \bar{\mu}_5 (2 \ln 2 - 1) + \cdots,$$

Plus tons of plots for arbitrary ω , q and μ , μ_5

$\text{Re}(\sigma_\chi), \kappa\bar{\mu}_5 = 0.25, \kappa\bar{\mu} = 0$




$\text{Im}(\sigma_\chi), \kappa\bar{\mu}_5 = 0.25, \kappa\bar{\mu} = 0$




$q = 0$

H.-U. Yee, (2009)

$\text{Re}(\sigma_\chi), \kappa\bar{\mu}_5 = 0.125, \kappa\bar{\mu} = 0.125$



$\text{Im}(\sigma_\chi), \kappa\bar{\mu}_5 = 0.125, \kappa\bar{\mu} = 0.125$



II) Non-linear corrections induced by the magnetic field

$$\vec{B} = \vec{B}(\vec{x}) \neq \vec{B}(t)$$

$$\mathcal{D}_0 = \frac{1}{2} - 18(2 \log 2 - 1) \kappa^2 \mathcal{B}^2$$

The correction is negative! Violates low bounds P. Kovtun and A. Ritz, (2008)

$$\omega = \left[\mp 1 + 36(1 - 2 \log 2) \kappa^2 \mathcal{B}^2 \right] 6\kappa \vec{q} \cdot \vec{B} - \left[\frac{1}{2} + 18(1 - 2 \log 2) \kappa^2 \mathcal{B}^2 \right] i q^2 - \frac{i}{8} q^4 \log 2 + \dots$$

The first term represents the chiral magnetic wave (CMW) D. E. Kharzeev and H.-U. Yee, (2011)

We see nonlinear in \mathcal{B} corrections to both the speed of CMW and its decay rate.
Also the degeneracy in the speed of the positive and negative modes is removed.

III) Non-linear transport induced by constant external e/m fields

The external fields are assumed to have constant background values $\vec{E}, \vec{B}, \vec{E}^a, \vec{B}^a$

$$\begin{aligned}\vec{\mathcal{E}}(\mathbf{x}_\alpha) &= \vec{E} + \epsilon \delta \vec{E}(\mathbf{x}_\alpha), & \vec{\mathcal{B}}(\mathbf{x}_\alpha) &= \vec{B} + \epsilon \delta \vec{B}(\mathbf{x}_\alpha), \\ \vec{\mathcal{E}}^a(\mathbf{x}_\alpha) &= \vec{E}^a + \epsilon \delta \vec{E}^a(\mathbf{x}_\alpha), & \vec{\mathcal{B}}^a(\mathbf{x}_\alpha) &= \vec{B}^a + \epsilon \delta \vec{E}^a(\mathbf{x}_\alpha).\end{aligned}$$

- Constant background $\vec{B} \neq 0, \quad \vec{E} = \vec{B}^a = \vec{E}^a = 0$

$$\mathbf{J}_{(0)}^t = \bar{\rho}, \quad \vec{\mathbf{J}}_{(0)} = 12\kappa\mu_5(\mathbf{B}) \vec{B} \quad \mathbf{J}_{5(0)}^t = \bar{\rho}_5, \quad \vec{\mathbf{J}}_{5(0)} = 12\kappa\mu(\mathbf{B}) \vec{B}$$

Index $_{(0)}$ means zeroth order in gradient expansion ($\epsilon = 0$)

CME is exact to all orders in \vec{B} . D. E. Kharzeev and H.-U. Yee (2011); A. V. Sadofyev and M. V. Isachenkov (2011), U. Gursoy and A. Jansen (2014); U. Gursoy and J. Tarrio (2015)
but there are gradient and \vec{E} -field corrections

- Constant background e/m fields \vec{B} & \vec{E} , weak field expansion, $\vec{B}^a = \vec{E}^a = 0$

$$\vec{J}_0 = \vec{E} + 12\kappa\mu_5(\mathbf{B}, \mathbf{E})\vec{B} + 72 \log 2 \kappa^2 \mu(\mathbf{B})\vec{E} \times \vec{B} - 36\pi^2 \kappa^3 \mu_5(\mathbf{B}, \mathbf{E}) (\vec{B} \times \vec{E}) \times \vec{E} + \dots,$$

$$\vec{J}_{5(0)} = 12\kappa\mu(\mathbf{B})\vec{B} + 72 \log 2 \kappa^2 \mu_5(\mathbf{B}, \mathbf{E})\vec{B} \times \vec{E} - 36\pi^2 \kappa^3 \mu(\mathbf{B}) (\vec{B} \times \vec{E}) \times \vec{E} + \dots,$$

Field-dependent chemical potentials

$$\mu(\mathbf{B}) = \frac{1}{2}\bar{\rho} + 18 (1 - 2 \log 2) \kappa^2 \bar{\rho} \mathbf{B}^2 + \dots,$$

$$\mu_5(\mathbf{B}, \mathbf{E}) = \frac{1}{2}\bar{\rho}_5 + 18 (1 - 2 \log 2) \kappa^2 \bar{\rho}_5 \mathbf{B}^2 + \frac{1}{8} (-\pi + 2 \log 2) 12\kappa \vec{B} \cdot \vec{E} + \dots.$$

- μ_5 is induced even in totally neutral plasmas $\bar{\rho} = \bar{\rho}_5 = 0$, via $(\vec{E} \cdot \vec{B})$.

- $\mathbf{B}^2 \vec{B}$ & $(\vec{B} \vec{E}) \vec{B}$ terms are the first nonlinear effects in CME
important for discussions of strong magnetic fields

$$\vec{J}_0 = \vec{E} + 12\kappa\mu_5(\mathbf{B}, \mathbf{E})\vec{B} + 72 \log 2 \kappa^2 \mu(\mathbf{B})\vec{E} \times \vec{B} - 36\pi^2 \kappa^3 \mu_5(\mathbf{B}, \mathbf{E}) (\vec{B} \times \vec{E}) \times \vec{E} + \dots,$$

$$\vec{J}_{5(0)} = 12\kappa\mu(\mathbf{B})\vec{B} + 72 \log 2 \kappa^2 \mu_5(\mathbf{B}, \mathbf{E})\vec{B} \times \vec{E} - 36\pi^2 \kappa^3 \mu(\mathbf{B}) (\vec{B} \times \vec{E}) \times \vec{E} + \dots,$$

- $(\vec{B} \times \vec{E})$ term leads to anomaly induced Hall current
(chiral Hall effect S. Pu, S.-Y. Wu, and D.-L. Yang, (2015))
- $\vec{J}_0 \sim (\vec{B} \times \vec{E}) \times \vec{E} = -E^2 \vec{B}$ (CME) + $(\vec{E} \cdot \vec{B})\vec{E}$ (CEE)
(χ KT E. V. Gorbar, I. A. Shovkovy, S. Vilchinskii, I. Rudenok, A. Boyarsky, and O. Ruchayskiy, (2016))

CEE Chiral Electric Effect Y. Neiman and Y. Oz, (2011)

- $\vec{J}_{5(0)} \sim (\vec{B} \times \vec{E}) \times \vec{E} = -E^2 \vec{B}$ (CSE) + $(\vec{E} \cdot \vec{B})\vec{E}$ (CESE)

CESE Chiral Electric Separation Effect X.-G. Huang and J. Liao, (2013), even when $\mu_5 = 0$

Towards all order non-linear constitutive relations

Linear in constant background times linear in inhomogeneous field perturbations

$$\begin{aligned}
 & \delta\rho_5 \vec{\mathbf{B}}, \quad \delta\rho \vec{\mathbf{B}}^a, \quad \left(\vec{\nabla} \cdot \delta\vec{\mathbf{E}}^a \right) \vec{\mathbf{B}}, \quad \left(\vec{\nabla} \cdot \delta\vec{\mathbf{E}} \right) \vec{\mathbf{B}}^a, \quad \vec{\mathbf{E}}^a \times \delta\vec{\mathbf{E}}, \quad \vec{\mathbf{E}}^a \times \vec{\nabla} \delta\rho, \\
 & \quad \vec{\mathbf{E}}^a \times \left(\vec{\nabla} \times \delta\vec{\mathbf{B}} \right), \quad \vec{\mathbf{E}}^a \times \delta\vec{\mathbf{B}}, \quad \vec{\mathbf{E}}^a \times \left(\vec{\nabla} \times \delta\vec{\mathbf{B}}^a \right), \quad \vec{\mathbf{E}}^a \times \delta\vec{\mathbf{B}}^a, \\
 & \quad \vec{\mathbf{E}}^a \times \vec{\nabla} \left(\vec{\nabla} \cdot \delta\vec{\mathbf{E}} \right), \quad \left(\bar{\rho} \vec{\mathbf{B}} + \bar{\rho}_5 \vec{\mathbf{B}}^a \right) \times \delta\vec{\mathbf{E}}, \quad \left(\bar{\rho} \vec{\mathbf{B}} + \bar{\rho}_5 \vec{\mathbf{B}}^a \right) \times \vec{\nabla} \delta\rho, \\
 & \left(\bar{\rho} \vec{\mathbf{B}} + \bar{\rho}_5 \vec{\mathbf{B}}^a \right) \times \left(\vec{\nabla} \times \delta\vec{\mathbf{B}} \right), \quad \left(\bar{\rho} \vec{\mathbf{B}} + \bar{\rho}_5 \vec{\mathbf{B}}^a \right) \times \delta\vec{\mathbf{B}}, \quad \left(\bar{\rho} \vec{\mathbf{B}} + \bar{\rho}_5 \vec{\mathbf{B}}^a \right) \times \left(\vec{\nabla} \times \delta\vec{\mathbf{B}}^a \right), \\
 & \quad \left(\bar{\rho} \vec{\mathbf{B}} + \bar{\rho}_5 \vec{\mathbf{B}}^a \right) \times \delta\vec{\mathbf{B}}^a, \quad \left(\bar{\rho} \vec{\mathbf{B}} + \bar{\rho}_5 \vec{\mathbf{B}}^a \right) \times \vec{\nabla} \left(\vec{\nabla} \cdot \delta\vec{\mathbf{E}} \right), \quad \vec{\mathbf{E}} \times \delta\vec{\mathbf{E}}^a, \quad \vec{\mathbf{E}} \times \vec{\nabla} \delta\rho_5, \\
 & \quad \vec{\mathbf{E}} \times \left(\vec{\nabla} \times \delta\vec{\mathbf{B}}^a \right), \quad \vec{\mathbf{E}} \times \delta\vec{\mathbf{B}}^a, \quad \vec{\mathbf{E}} \times \left(\vec{\nabla} \times \delta\vec{\mathbf{B}} \right), \quad \vec{\mathbf{E}} \times \delta\vec{\mathbf{B}}, \quad \vec{\mathbf{E}} \times \vec{\nabla} \left(\vec{\nabla} \cdot \delta\vec{\mathbf{E}}^a \right), \\
 & \quad \left(\bar{\rho}_5 \vec{\mathbf{B}} + \bar{\rho} \vec{\mathbf{B}}^a \right) \times \delta\vec{\mathbf{E}}^a, \quad \left(\bar{\rho}_5 \vec{\mathbf{B}} + \bar{\rho} \vec{\mathbf{B}}^a \right) \times \vec{\nabla} \delta\rho_5, \quad \left(\bar{\rho}_5 \vec{\mathbf{B}} + \bar{\rho} \vec{\mathbf{B}}^a \right) \times \left(\vec{\nabla} \times \delta\vec{\mathbf{B}}^a \right), \\
 & \quad \left(\bar{\rho}_5 \vec{\mathbf{B}} + \bar{\rho} \vec{\mathbf{B}}^a \right) \times \delta\vec{\mathbf{B}}^a, \quad \left(\bar{\rho}_5 \vec{\mathbf{B}} + \bar{\rho} \vec{\mathbf{B}}^a \right) \times \left(\vec{\nabla} \times \delta\vec{\mathbf{B}} \right), \quad \left(\bar{\rho}_5 \vec{\mathbf{B}} + \bar{\rho} \vec{\mathbf{B}}^a \right) \times \delta\vec{\mathbf{B}}, \\
 & \quad \left(\bar{\rho}_5 \vec{\mathbf{B}} + \bar{\rho} \vec{\mathbf{B}}^a \right) \times \vec{\nabla} \left(\vec{\nabla} \cdot \delta\vec{\mathbf{E}}^a \right).
 \end{aligned}$$

The first terms $\delta\rho_5 \vec{\mathbf{B}}, \delta\rho \vec{\mathbf{B}}$ are responsible for the chiral magnetic wave (CMW)

IV) Constant magnetic and time-dependent electric fields

$$\vec{B} = \vec{B}, \quad \vec{E} = \vec{E}(t) \neq \vec{E}(\vec{x}), \quad \vec{B}^a = \vec{E}^a = 0, \quad \bar{\rho} = \bar{\rho}_5 = 0$$

$$\partial_\mu \mathbf{J}^\mu = 0, \quad \partial_\mu \mathbf{J}_5^\mu = 12\kappa \vec{E} \cdot \vec{B} \longrightarrow \mu_5 \neq 0$$

• Gradient expansion

$$\begin{aligned} \vec{J} = 12\kappa\mu_5\vec{B} + \vec{E} - \frac{\log 2}{2}\partial_t\vec{E} - \frac{\pi^2}{24}\partial_t^2\vec{E} - \left(\frac{3}{2}\pi + 3\log 2\right)\kappa\partial_t\rho_5\vec{B} \\ + 9\pi^2\kappa^3\rho_5\left(\vec{B} \times \vec{E}\right) \times \vec{E} + 12\#_1\kappa\partial_t^2\rho_5\vec{B} + \mathcal{O}\left(\partial^4\right), \end{aligned}$$

$$\begin{aligned} \vec{J}_5 = 12\kappa\mu\vec{B} - 36\log 2\kappa^2\rho_5\vec{B} \times \vec{E} + \frac{3}{2}\left(\pi^2 + 3\pi\log 2 + 6\log^2 2\right)\kappa^2\partial_t\rho_5\vec{B} \times \vec{E} \\ - \frac{3}{8}\left(48\mathcal{C} + \pi^2 - 12\pi\log 2\right)\kappa^2\rho_5\vec{B} \times \partial_t\vec{E} + \mathcal{O}\left(\partial^4\right), \end{aligned}$$

where \mathcal{C} is a Catalan constant and $\#_1$ is known numerically only $\#_1 \approx 0.362$.

$$\mu = 0 + \mathcal{O}\left(\partial^3\right), \quad \mu_5 = \frac{1}{2}\rho_5 + \frac{3}{2}(\pi - 2\log 2)\kappa\vec{E} \cdot \vec{B} + 18(1 - 2\log 2)\kappa^2\rho_5\mathbf{B}^2 + \mathcal{O}\left(\partial^3\right).$$

- Weak electric field expansion

$$\rho_5 \sim \mathcal{O}(\epsilon), \quad \vec{\mathcal{E}}(\mathbf{t}) \sim \mathcal{O}(\epsilon), \quad \vec{\mathbf{B}} \sim \mathcal{O}(\epsilon^0)$$







Linear in ϵ





$$\mathbf{J}^t = 0, \quad \vec{\mathbf{J}} = \sigma_e [\partial_t] \vec{\mathcal{E}} + \kappa \tau_1 [\partial_t] \rho_5 \vec{\mathbf{B}} + \kappa^2 \tau_2 [\partial_t] (\vec{\mathcal{E}} \cdot \vec{\mathbf{B}}) \vec{\mathbf{B}}; \quad \mathbf{J}_5^t = \rho_5, \quad \vec{\mathbf{J}}_5 = 0$$

The electric current is put on-shell,





$$\mathbf{J}^i = \sigma_{ij} \mathcal{E}_j, \quad \sigma_{ij} = \underbrace{\sigma_e}_{\sigma_T} \left(\delta_{ij} - \frac{\mathbf{B}_i \mathbf{B}_j}{\mathbf{B}^2} \right) + \underbrace{\left[\sigma_e - \left(\frac{12}{i\omega} \tau_1 - \tau_2 \right) \kappa^2 \mathbf{B}^2 \right]}_{\sigma_L} \frac{\mathbf{B}_i \mathbf{B}_j}{\mathbf{B}^2},$$

In DC limit τ_1^0 is known analytically K. Landsteiner, Y. Liu, and Y.-W. Sun, (2015);
M. Ammon, S. Griener, A. Jimenez-Alba, R. P. Macedo, and L. Melgar, (2016)





To enhance the effect of anomaly, $\delta G^{\mathbf{L}} = G^{\mathbf{L}} - G^{\mathbf{T}}$



Conclusions

- An off-shell constitutive relation for $U(1)$ current consists of a momenta-dependent diffusion term and two conductivities. Certain universality between dissipative transport coefficients η and \mathcal{D} is observed.
- Causality restoration: at large momenta, the effective diffusion TCF is a decreasing function of both frequency; the corresponding memory function has support in the past only.
- We have re-examined transport coefficients induced by the chiral anomaly. We seem to be able to rediscover all known anomaly-induced effects within one and the same holographic model, without introducing any additional inputs or model assumptions, which appeared in the literature
- For linearized problem, we have completely determined all anomaly-induced TCFs to all orders in the gradient expansion
- For nonlinear problem with constant external fields, we have found E -induced corrections to CME/CSE, and B -induced modifications to CMW, and diffusion coefficient D
- There seem to be enhancements of anomaly-induced affects at finite frequencies